



PRELIMINARY PHYTOCHEMICAL SCREENING AND ANTIMICROBIAL EFFECT TOWARD *VIBRIO* *PARAHAEMOLYTICUS* ISOLATED FROM WHITE LEG SHRIMP (*PENAEUS VANNAMEI*) OF THE AQUEOUS HERBAL EXTRACT OF *WEDELIA CHINENSIS*

Tran Nguyen Ngoc^{1*}, Nguyen Ngoc Phuoc¹, Nguyen Quang Linh²

Abstract

The trend of using natural compounds or herbs containing antibacterial characteristics for disease prevention and treatment is becoming common, because of its environmentally friendly, and ensures food safety and animal harm reduction. There are numerous Vietnamese herbs that contain antibacterial and environmentally friendly features, however, the use of herbs in aquaculture is still limited. This study aims to determine the antibacterial components of *Wedelia chinensis* extract for the prevention of diseases caused by *Vibrio parahaemolyticus* in white-leg shrimp (*Penaeus vannamei*). There were 36 samples of grass (*W. chinensis*), which were collected in Central Vietnam and were extracted in 3 different solvents (distilled water, methanol 99.8%, and ethanol 96%) for phytochemical screening by the standard phytochemical assays. The results of the preliminary phytochemical screening in this study showed the extract in 3 different solvents containing various chemical compounds such as alkaloids, saponin, tannin, flavonoids, carbohydrates, organic acids, fixed oil, and fats. The total polyphenols and flavonoids of *W. chinensis* extract were analyzed by folin-ciocalteau reagent (for polyphenols) and by $AlCl_3$ (for flavonoids). The highest amount of total polyphenols and flavonoids were recorded in the *W. chinensis* extract in the solvent of methanol 99.8% (74.33 ± 4.49 mg GAE/g and 24.59 ± 2.19 mg RE/g dry weight, respectively), followed by in the solvent of ethanol 96% (73.65 ± 5.44 mg GAE/g and 20.63 ± 4.30 RE/g of dry weight respectively). The lowest of total polyphenols and flavonoids was observed in the solvent of distilled water (53.07 ± 1.48 mg GAE/g and 3.20 ± 0.07 mg RE/g, respectively). The efficiency of aqueous herbal extraction is highest in the solvent of distilled water (up to 18.2%), followed by in the solvent of ethanol 96% (11.3%), and in the solvent of methanol 99.8%, (11%). The results of antagonistic to *V. parahaemolyticus* testing of *W. chinensis* extract in 3 solvents showed that all *W. chinensis* extract inhibited the growth of *V. parahaemolyticus*, in which the diameter of the antibacterial zone of *W. chinensis* extracts in methanol solvent was highest up to 14.7 ± 0.91 mm. The minimum inhibitory concentrations (MIC) of *W. chinensis* extract in the solvents: distilled water, methanol 99.8%, and ethanol 96% against *V. parahaemolyticus* were 2500 mg/L, 15.5 mg/L to 15.5 mg/L, respectively. The minimum bactericidal concentration (MBC) of *W. chinensis* extracts was 31.25 mg/L in the solvent of methanol 99.8%, and 62.5 mg/L in the solvent of distilled water. Toxicity test of *W. chinensis* extract in methanol solvent 99.8% in white leg shrimp (*P. vannamei*) showed the concentration of 31.25 mg/L (MBC) was safe for shrimp.

Keywords: *W. chinensis*, *V. parahaemolyticus*, solvents, quantification, and qualification.

^{1*}Faculty of Fisheries, University of Agriculture and Forestry, Hue University, 102 Phung Hung St. Hue 49000, Vietnam

²Faculty of Animal Sciences and Veterinary Medicine, University of Agriculture and Forestry, Hue University, 102 Phung Hung St. Hue 49000, Vietnam

*Corresponding Author: Tran Nguyen Ngoc, ORCID: 0000-0003-4867-9596

*Faculty of Fisheries, University of Agriculture and Forestry, Hue University, 102 Phung Hung St. Hue 49000, Vietnam

Other authors: Nguyen Quang Linh, E-mail nguyen.linh@huaf.edu.vn or nguyenquanglinh@hueuni.edu.vn

ORCID: 0000-0002-8424-1148

Nguyen Ngoc Phuoc, nguyenngocphuoc@huaf.edu.vn,

ORCID: 0000-0003-1928-7462

DOI: 10.53555/ecb/2022.11.12.204

Introduction

The trend of using natural compounds or herbs containing antibacterial biological activities for disease prevention and treatment in aquaculture is being common because of its environmentally friendly, and ensuring food safety (Cos et al 2006) [22]. Natural compounds such as herbs are becoming increasingly common alternative antibiotics to prevent bacterial disease in aquaculture and ensure food safety. There are numerous Vietnamese herbs which contain antibacterial and environmentally friendly features. Several studies have shown the important role of herbs in the competitive exclusion of pathogenic bacteria, providing nutrient addition, and enzymes that contribute to fish digestion, enhancement of the fish's immune response, and disease resistance [26]. The extract of gallic could be used as an antibacterial compound for shrimp [11]. Nguyen Ngoc Phuoc et al (2007) successfully tested the antifungal ability of betel leaf extract in five species of *Saprolegnia* and *Aphanomyces* infected in fish [10]. The studies of Nguyen Quang Linh (2010) and Pham Thi Hai Yen et al (2019) [6],[13] showed the antibacterial effect of betel leaf extract and *Phyllanthus amarus* extract on pathogenic *Vibrio* strains isolated in shrimp and red drum fish in Vietnam. However, the information on phytochemical and antimicrobial compound in herbs are still limited. This study aimed to preliminary screen the phytochemical compounds and to determine the antimicrobial effect of the aqueous herbal extract of *W. chinensis* toward *V. parahaemolyticus* isolated from white leg shrimp (*P. vannamei*) and to identify the toxicity of this extract in the shrimp (*in vivo* test) to providing the scientific information for using this herbal extraction in the prevention of bacterial disease in shrimp.

2. Materials and methods

2.1. Materials, Microorganisms and animals

W. chinensis (root, stem, and leaf) was collected from Thua Thien Hue province, Vietnam. The grass parts were shade-dried in an oven (FSV 280, China) at 60°C for 8 hours and ground to obtain moderately coarse powder in the machine (Phillips HR 3573/90, China) and were extracted in 3

different solvents: distilled water, methanol 99.8% and ethanol 96% according to the method of Trieu Thi Thanh Hang et al (2018) [2]

Bacterial strain *V. parahaemolyticus* (NNP 001) (Figure 1) was isolated from white leg shrimp that was infected Acute hepatopancreas Necrosis Disease (AHPND cultured in Thua Thien Hue, was storage in Glycerol 15% at -20 °C and supplied from the Laboratory of Fish Pathology, the Faculty of Fishery, University of Agriculture and Forestry, Hue University, Hue city, Vietnam [9].

V. parahaemolyticus NNP 001 was firstly recovered on Tryptic Soy Agar (TSA, Himedia, India) supplemented with 2% NaCl, then sub-cultured in Tryptic Soy Broth (TSB, Himedia, India) supplemented with 2% NaCl and incubated at 30 °C and 180 rpm for 24 h in the GFL 3032 incubator (GFL, Germany).

Bacterial density after culture was determined by optical density (OD) = 1 at 600 nm on UV-VIS spectrophotometer (U2900, Hitachi, Japan) (equivalent to a bacterial density of 10⁹ CFU/mL, based on the standard curve for *V. parahaemolyticus* strain that provided by Laboratory of Fish Pathology, Faculty of Fisheries, University of Agriculture and Forestry, Hue University), then diluted to 10⁶ CFU/mL to test the antibacterial ability of the *W. chinensis* extract in different solvents toward *V. parahaemolyticus*.

Juvenile *P. vannamei* weighing 1.5 - 2.0 g were purchased from Phan Toan hatchery at Phu Thuan Commune, Phu Vang district, Thua Thien Hue province, Vietnam. Juvenile *P. vannamei* was transferred by air-conditioned car to the wet lab of the Faculty of Fishery, University of Agriculture and Forestry, Hue University. The shrimp were maintained in 2 000 L fiber glass tanks at 28 °C ± 2 °C and four times a day fed the basal diet for 14 days in the aquaria, before use. Shrimp used in this study were quarantined for free Acute Hepatopancreatic Necrosis Disease (AHPND), White spot disease (WSD), and Yellow head disease (YHD) at the Veterinary Clinic, Sub-Department of Animal Husbandry and Veterinary Medicine in Thua Thien Hue (Figure 2).




Figure 1. *Vibrio parahaemolyticus*



Figure 2. Experimental white leg shrimp




Figure 3. The extraction process of *W. chinensis* extract

2.2. Variables

- The phytochemical compounds and antimicrobial effect of the aqueous herbal extract of *W. chinensis* toward *V. parahaemolyticus*
- The minimum inhibitory concentration (MIC) and the minimum bactericidal concentration (MBC) of the *W. chinensis* extract toward *V. parahaemolyticus*.
- The toxicity of the *W. chinensis* extract in white-leg shrimp *P. vannamei*

2.3. Methods

- Herbal extraction in different solvent was carried out according to Trieu Thi Thanh Hang et al (2018) [2], briefly 10g of *W. chinensis* powder were extracted separately with 100 mL distilled water, or 100 mL methanol 99.8%, or 100-mL ethanol 96% in a water bath 100 °C for 20-30 minutes and leave at room temperature for 24 h. The extracts were processed by ultrasound for 1 h then filtered and concentrated to dry in a rotary evaporator for eliminating the solvents. The semi-solid extract of leaf, stem, and root was thus obtained, and was re-dissolved in water and stored in a refrigerator for phytochemical screening and antimicrobial test (Figure 3) [2].

- The method of phytochemical screening of *W. chinensis* extract in different solvents (methanol 99.8%, ethanol 96%, or distilled water) was conducted according to the method of Tran Hung and Nguyen Viet Kinh (2015) [3]
- The total polyphenol of the initial extract was determined according to the Folin-Ciocalteu (a mixture of sodium wolframite and sodium molybdate) method. If phenol was presented in the extract, an oxidation-reduction reaction would have occurred. The – OH groups in phenolic will be converted into quinol groups to a complex form of blue color with a wide spectrum of light absorption. The sample was tested with a standard for reaction with the Folin-Ciocalteu reagent; then, the optical density was measured at the maximum absorption wavelength. The reaction proceeded in condition avoiding light. Formulate the standard curve of gallic acid in the appropriate concentration range. The total polyphenol content was calculated using the standard curve of gallic acid [16], [29].
- Defined the standard curve of gallic acid Preparation of the base standard solution and diluted gallic acid titration solutions: Weigh exactly 10.0 mg of gallic acid into rated vessels

- with a capacity of 100 mL, dissolve, and add to the mark of 10mL with methanol. A standard solution of gallic acid at a concentration of 1 mg/mL was obtained. From this solution, dilute with methanol to standard gallic acid with a concentration of 50; 75; 100; 125; 150; 175 µg/mL (marked C1 to C6 respectively) [16], [29].
- Calibration step: Conduct a reaction between 0.2 mL of each diluted solution (C1 to C6), 0.8 mL of distilled water, and 1 mL of 10% Folin-Ciocalteu reagent, shake well. Leave the reaction for 5 minutes. After that 2.5 mL of Na₂CO₃ 7.5%. Measure the absorption at OD_{760nm} after 30 minutes at room temperature. From the measured optical density value, build a gallic acid benchmark.
 - Sample preparation: Weight 20.0 mg of extracts and dissolve these extracts with methanol in a 10 mL rated vessel for suitable test concentration. Conduct a reaction with 10% Folin-Ciocalteu reagent similar to the above. From the obtained optical density value and based on the results of the linearity between the concentration and absorption of gallic acid standard solution, the total polyphenol content in each herbal extract was determined.
 - Control sample included 0.2 mL methanol, and 1.8 mL distilled water, and 2.5 mL Na₂CO₃. The total polyphenol content (TPC) was calculated in mg of gallic acid equivalent to 1g of herbal extract (mg GAE/g) according to the formula:

$$\text{TPC (mg GAE/g extract)} = \frac{C_x \times V \times K}{m_o}$$

Where:

TPC was the total polyphenols present in 1 g extract (mg GAE/g extract)

C_x was the concentration of total polyphenols in the photogrammetric solution in terms of gallic acid (mg/mL)

V was the initial volume of the extraction solution (mL)

K was the dilution factor

m_o was the initial weight of extract (g)

Determine the total flavonoid in the *W. chinensis* extract was followed the colorimetric method using AlCl₃ [28], [33] based on the standard curve of rutin

- Accurately weigh 10.0 mg of rutin into a rated vessel with a capacity of 100 mL, dissolve with methanol solvent, and rate to the mark of 10 mL to get a standard rutin solution with concentration C_o=1 mg/mL. Took exactly 1 mL

of this solution with a micropipette, put it in a 10 mL rated vessel, and diluted it to the mark with methanol, obtaining a standard rutin solution with a concentration of C_x=100 µg/mL. This solution with methanol was into a series of diluted solutions C1-C6, whose concentration was 20.0; 27.5; 35.0; 42.5; 50.0; and 57.5 µg/mL [28], [33].

- Calibration: Put 2 mL of each diluted standard solution (C1-C6) in test tubes and 2 mL AlCl₃ 2% shake well. For the reaction to take place for 10 minutes at room temperature, measure absorption at wavelength 430 nm, using a white sample of 2 mL of rutin standard at the corresponding concentration and 2 mL of methanol. Construct a rutin calibration based on the obtained value of optical density.
- Test sample preparation: Accurately weigh 10.0 mg extract in methanol or ethanol and 70.0 mg extract in distilled water, dissolved in methanol solvent in a 10 mL rated vessel for appropriate concentration. The reaction with the 2% AlCl₃ reagent as above, using 2 mL control sample, and 2 mL methanol. From the obtained optical density value and the results of the linear concentration and absorption of the rutin standard, the total flavonoid content of each extract was determined [28], [33]. The total flavonoid content (TFC) was calculated in mg of rutin that equivalent to 1g of herb (mg RE/g extract) according to the formula:

$$\text{TFC (mg RE/g extract)} = \frac{C_x \times V \times K}{m_o}$$

Where: TFC was the total amount of flavonoids present in 1 g of dry herb (mg RE/g extract) and C_x was the total flavonoid in rutin solution measured by colorimetric (mg/mL)

- The efficiency of extraction in different solvents was calculated as follows:

$$H = \frac{M_{cao}}{M_{k\grave{h}o}} \times 100$$

Where: H: extract yield (%); M_{cao}: Total weight of extract (g); M_{k_ho}: total weight of dried herb (g)

- Evaluating the antimicrobial effect of *W. chinensis* extract toward *V. parahaemolyticus* which caused the acute hepatopancreatic necrosis disease in white-leg shrimp

The antimicrobial effect of *W. chinensis* extract in different solvents followed the method of

Tucker et al (2009) [35] including the following steps:

- Spread the bacteria evenly over the prepared medium of TSA+ 2% NaCl, and allow to dry for 3–5 minutes at room temperature before making a 6 mm diameter well at appropriate spacing on the bacteria-spread agar plate.
- Accurately put 50 µl of distilled water into a well (1) for negative control.
- Put 50 µL of *W. chinensis* extract in 99.8% methanol into a well (2) (concentration equivalent to 2500 µg/well).
- Put 50 µL of *W. chinensis* extract in 96% ethanol into a well (3) (concentration equivalent to 2500 µg/well).

- Put 50 µL of *W. chinensis* extract in distilled water into a well (4) (concentration equivalent to 2500 µg/well).
- Placed an antibiotic disc of amoxicillin (125 µg) (positive control) on the plate.

Then allow 15 minutes for the extract to diffuse into the agar layer. Check the diameter of the antimicrobial zones after 24 h at 28–30 °C. The degree of resistance to *V. parahaemolyticus* bacteria of the *W. chinensis* extract was determined according to Faikoh et al (2014) [25] (Table 1).

Table 1. Antimicrobial activity of *W. chinensis* extract (Faikoh et al 2014)

Diameter of an antimicrobial zone (mm)	Inference
≥ 15	Strong
7.5 – 15	Medium
< 7.5	Weak
0	Resistance

- The minimum inhibitory concentration (MIC) of each extract was performed on the 96-well dish according to the dilution method of CLSI (2012) [21]. The minimum bactericidal concentration (MBC) was the lowest concentration in the range of extracts that can kill all bacteria in the well (bacteria that do not grow when re-subculture the solution in these wells on TSA plates) (Figure 8) [20].

- Toxicity test

The experiment was designed in plastic tanks (V = 100L) sterilized with chlorine 100 ppm and dried. Then supply seawater (salinity 30 ‰) to about 2/3 of the tank volume and continuous aeration. White leg shrimp was distributed with a density of 30 shrimp per tank. Before conducting the experiments, shrimp were tested for free *Vibrio* by taking hepatopancreatic samples of 5 shrimp and spread directly on TCBS agar plates, observing bacterial growth after 24 hours at 28 °C (Figure 4). The experimental design included:



Figure 4. Experimental tanks

Experimental trials with 3 different extraction concentrations (treatment) and 1 control test. Each treatment was performed in triplicates. Where:

Treatment (NT) 1: Shrimp were fed commercial feed added *W. chinensis* extract at MBC concentration.

Treatment (NT) 2: Shrimp were fed commercial feed added *W. chinensis* extract at 20x MBC concentration.

Treatment (NT) 3: Shrimp were fed commercial feed added *W. chinensis* extract at 40x MBC concentration.

Shrimp were fed commercial feed with or without *W. chinensis* extract 3 times a day at 6h, 12h, and 18h. The feed was provided at 2.0 % of the body weight of the total shrimp in each tank. Monitoring and evaluation of survival for 7 days.

Based on the cumulative mortality, the LD₅₀ (Lethal dose) was calculated according to the formula of Reed & Muench (1938) [34].

2.4. Data analysis

The data was processed using Excel 2016 and SPSS 16.0 software, a one-factor ANOVA variance to compare differences in antimicrobial ring diameters. Statistical testing was done at a significance level of $P \leq 0.05$, by LSD test.

Table 2. Biological substances present in the extracts (following the method of Deepa and Padmaja 2014 with slightly modified)

Test for	Reagents	Reaction	Inference		
			Methanol	Ethanol	Distilled Water
Fats	Extract pressed between filter paper	Oil stain develops	+	+	-
Fixed oils	Extract pressed between filter paper	Oil stain develops	+	+	-
Carotenoids	H ₂ SO ₄	Blue color	+	+	+
Alkaloids	Iodine in water	Reddish-brown precipitate	+	+	+
Flavonoid-pyron	HCl	Pink or red color	+	+	+
Tannins	10% alcoholic ferric chloride	Dark blue or greenish grey	-	+	+
Saponin	Distilled water	A 1cm layer of foam	-	-	+
Organic acids	Na ₂ CO ₃	Gas released.	+	+	+
Carbohydrates	Copper sulphate+ potassium sodium tartarate+NaOH	Reddish-brown precipitate	+	+	+
Flavonoid-Proanthocyanidin	KOH 10%	Dark blue	+	+	+

Note: (+): present; (-): absent

Table 3. Composition of polyphenol and flavonoid in *W. chinensis* extract in 3 solvents

<i>W. chinensis</i> extract in solvents	Polyphenol (mg GAE/g)	Flavonoid (mg RE/g)
Methanol	74.33 ± 4.49 ^b	24.59 ± 2.19 ^b
Ethanol	73.65 ± 5.44 ^b	20.63 ± 4.30 ^b
Distilled water	53.07 ± 1.48 ^a	3.20 ± 0.07 ^a

Note: a, b the differences same column significantly, $P < 0.05$

Table 4. The efficiency of *W. chinensis* extract

<i>W. chinensis</i> extract in solvents	Sample weight (g)	The weight of extract (g)	Efficiency (%)
Methanol	10	1.1	11
Ethanol	10	1.13	11.3
Distilled water	10	1.82	18.2

Table 5. Diameter of antimicrobial zone of *W. chinensis* extract (mm) in different solvents toward *V. parahaemolyticus*

<i>W. chinensis</i> extract in solvents	Diameter of the antibacterial zone (mm)
Methanol 99.8%	14.7 ^a ± 0.91
Ethanol 96%	12.7 ^b ± 1.92
Distilled water	12.1 ^b ± 1.59
Positive Control (Amoxicillin)	9.2 ^{a,b} ± 5.3
Negative Control (Distilled water only)	0 ^c

Note: a, b, c the differences in the same column significantly, $P < 0.05$

Table 6. Determination of the minimum inhibitory concentration (MIC) of the extract of the *W. chinensis* extract toward *V. parahaemolyticus*.

<i>W. chinensis</i> extract Dilution concentration (mg/L)	Bacterial concentration (CFU/mL)	Minimum Inhibitory Concentration (MIC)		
		Methanol	Distilled Water	Ethanol
5000		-	-	-
2500		-	-	-
1000		-	+	-
500		-	+	-
250	10 ⁶ CFU/mL	-	+	-
125		-	+	-
62.5		-	+	-
31.25		-	+	+
15.5		+	+	+

Note: (+): bacteria growth, (-): no bacteria growth

Table 7. Minimum inhibitory concentrations (MIC) and minimum bactericidal concentrations (MBC) of *W. chinensis* extract toward *V. parahaemolyticus*.

<i>W. chinensis</i> extract in solvents	Minimum Inhibitory Concentration (MIC) (mg/L)	Minimum Bactericidal Concentration (MBC) (mg/L)
Distilled water	2500	5000
Methanol	15.5	31.25
Ethanol	15.5	62.5




Figure 5. standard curve acid galic




Figure 6. standard curve rutin




Figure 7. Antimicrobial activities of *Wedelia chinensis* extract in different solvents (methanol, ethanol, Distilled water) toward *Vibrio parahaemolyticus*




Figure 8. MIC determination of *Wedelia chinensis* toward *Vibrio parahaemolyticus*




Figure 9. Mortality percentage of shrimp after 7 days of feeding with *Wedelia chinensis* extract




Figure 10. Clinical signs of moribund shrimp when fed with high dose of *W. chinensis* extract, (A) Hepatopancreas of shrimp was pale when fed with *W. chinensis* extract at a dose of 1250 mg/L (treatment 3), (B,C) healthy shrimp with dark hepatopancreas when the fed at a dose of 31.25 mg/L of *W. chinensis* (treatment control and treatment 1)

3. Results and discussions

3.1. Results of the biological compound composition of *W. chinensis*

- Qualitative determination of the biological component of *W. chinensis* extract

The qualitative determination of biological component screening of *W. chinensis* extract in the solvent of methanol, ethanol, or distilled water showed the presence of biochemical compounds had good antibacterial effects and helped to enhance the immune system's response and increased the survival rate for white leg shrimp when infected *V. parahaemolyticus*. The results also indicated that the *W. chinensis* extracts in all three solvents contained components such as alkaloid, flavonoid, carotenoid, fat, oil, tannin, organic acid, and reducing agents (Table 2). Especially, the flavonoid content was most abundant in the *W. chinensis* extracts in all three solvents. Saponin, meanwhile, was only present in aqueous solvents. The results of Govindappa et al (2011) on biological compounds in the stems and leaves of *W. trilobata* (L.) indicated the presence of phenolic, flavonoid, and alkaloid in these parts [26]. According to Tran Thanh Men et al (2019), when studying the ability to inhibit seed germination of the *W. trilobata* (L.) extract showed the presence of phenolic, tannin, flavonoid, and alkaloid were in all three parts of stem, leaf, and flower [8]. The presence of tannin in the extract of the *W. trilobata* leaves in ethanol solvent had an antibacterial effect. The antimicrobial activity of tannin was indicated by its ability of inhibiting extracellular enzymes of microorganism (Muhammad and Mudi (2011); Akiyama et al (2001); Doss et al (2009) [30] [15], [24]. Tannin also inhibited the growth of microorganism by

forming metal ion complexes or inhibiting phosphorylation. Ahn (2017) identified that alkaloids, glycosides, polyphenols, and terpene from plant extracts have antimicrobial abilities [14]. Therefore, the result of the qualitative determination of some biochemical compounds present in the *W. chinensis* extract helps to explain the antibacterial properties of the *W. chinensis* extracts toward *Vibrio* infected in white-leg shrimp. Many studies determined biological compounds from plants that were able to kill *Vibrio*. The study of Rahman and Rashid (2008) showed the compounds of eclalbasaponin I and eclalbasaponin II from ginseng (*Eclipta prostrata*) extract inhibited the growth of *V. parahaemolyticus* and *V. mimicus* at concentrations of 100 µg.

In addition, the study of Munaeni et al (2019) noted 7-hydroxy-1,2-dimethoxyxanthoneisoquinolinein, naphthalen, and phenol from the extract of *Eleutherine bulbosa* in ethanol solvent could inhibit the growth of *V. parahaemolyticus* by damaging structure of bacterial cell wall and reducing bacterial cells attached [31]. Furthermore, the herb extract also contained biochemical compounds such as flavonoid, alkaloid, tannin, phenol, oil, terpenoid, glycoside, and saponin, which provided many benefits to cultured species such as appetite stimulation, stress reduction, growth promotion, immune stimulation, anti-pathogen activity and promotion of its maturity [18], [19]. Thus, the results of this study showed the biological compound in the *W. chinensis* extracts in three solvents (methanol, or ethanol, or distilled water)

they have great potential applications in aquaculture.

3.2. The total of polyphenol and flavonoid in the extract of *W. chinensis* in different solvents

3.2.1. Defined the standard curve of gallic acid and rutin

To determine the polyphenol content in the *W. chinensis*, the standard curve of gallic acid was formulated with concentration levels of 50; 75; 100, 125; 150, and 175 $\mu\text{g/mL}$. The gallic acid calibration equation in methanol (Figure 5) showed that the optical density values obtained from different gallic acid concentrations exhibit a strong correlation ($R^2 = 0.9888$) and the linear equation obtained from gallic acid concentrations was $y = 0.0039x + 0.0441$ (Figure 5).

The standard curve of rutin was formulated with concentration levels of 20; 27.5; 35; 42.5; 50 and 57.5 $\mu\text{g/mL}$ to determine the flavonoid content in the *W. chinensis* extract (Figure 6). The study results showed the optical density values obtained from different rutin concentrations exhibit a strong correlation ($R^2 = 0.9659$) and the linear equation obtained from these rutin concentrations was $y = 0.0059x + 0.0563$ (Figure 6).

3.2.2. Total polyphenol and flavonoid in the *W. chinensis* extract

Flavonoid and polyphenol are two important groups of biochemical compounds that are highly bioactive and especially antimicrobial (Kumar et al (2011); Courts-Williamson, (2015) [27], [23]. Therefore, the quantity of total flavonoid and total polyphenol are two important indicators to evaluate the antimicrobial ability of herbal extract. The results of total flavonoid and polyphenol are presented in Table 3. The total polyphenol and flavonoid of *W. chinensis* extract were analyzed by folin-ciocalteau reagent (for polyphenol) and by AlCl_3 (for flavonoid). The highest amount of total polyphenol and flavonoid were recorded in the *W. chinensis* extract in the solvent of methanol 99.8% (74.33 ± 4.49 mg GAE/g and 24.59 ± 2.19 mg RE/g of extract respectively), followed by those in the solvent of ethanol 96% (73.65 ± 5.44 mg GAE/g and 20.63 ± 4.30 mg RE/g of extract respectively). The lowest total polyphenol and flavonoid was observed in the extract of *W. chinensis* in the solvent of distilled water (53.07 ± 1.48 mg GAE/g and 3.20 ± 0.07 mg RE/g, respectively). This result was similar to the previous study of Neelam et al (2012) when studying the biochemical compounds in the extract of *W. trilobata*. The total polyphenol content of 74.38 ± 1.03 mg GAE/g and a total

flavonoid of 16.67 mg RE/g were determined in the extract of this grass in the ethanol solvent [32]. In addition, the study of Tran Thanh Men et al (2019) [8], showed that the flavonoid content in the extract of stem, leaves, and flower of *W. trilobata* was 22.70%, 28.84%, 55.81%, and the total polyphenol content was 30.04%, 30.66%, and 50.62%, respectively. Thus, the polyphenol in the extract of *W. chinensis* in methanol or ethanol was similar to the extract of *W. trilobata* in ethanol, but the flavonoid content in the extract of *W. chinensis* in methanol or ethanol was higher than the extract of *W. trilobata* in ethanol, by contrast, the polyphenol and flavonoid in the extract of *W. chinensis* in distilled water were lower than those in the extract of *W. trilobata* in ethanol. According to the study of Trinh Thi Trang and Nguyen Thanh Hai (2016), when using different types of solvents for herbal extraction gave different dissolved biochemical compounds [12]. For example, using ethanol solvents for betel leaf extraction gave more biochemical compounds than using methanol and distilled water solvents. Thus, our research results show that highly polar solvents (ethanol and methanol) gave the highest ability to extract biochemical ingredients in the *W. chinensis* extract.

3.2.3. The efficacy of *W. chinensis* extraction in different solvents

The efficiency of herbal extraction was highest in the solvent of distilled water (up to 18.2%), followed by those in the solvent of ethanol 96% (11.3%), or in the solvent of methanol 99.8% (11%) (Table 4).

These results showed that different solvents used for extraction gave a difference in the efficacy of the extraction of herbs. According to the study of Turker et al (2009) [35], the efficiency of herbal extraction and the active ingredient obtained in the extract were different depending on the type of solvents used. In particular, using methanol or ethanol as a solvent for herbal extraction gave higher antibacterial capacity in both Gram-positive and Gram-negative bacteria than those in distilled water. In addition, these authors also believed that extracts in all three solvents have great potential in disease control in aquaculture. Methanol was a higher polar solvent than ethanol or distilled water, then herbal extraction in methanol gave a higher capability of natural compounds found than others. These biochemical compounds in herbs can protect aquatic animals from pathogenic microorganisms in water.

3.3. Antimicrobial effect of *W. chinensis* extract toward *V. parahaemolyticus*

The extract of *W. chinensis* in all 3 solvents showed strong antibacterial effects toward *V. parahaemolyticus* that caused acute hepatopancreatic necrosis on white-leg shrimp cultured in Thua Thien Hue (Figure 7). The diameter of the antimicrobial zone of *W. chinensis* extracts in methanol solvent was highest and higher than that in the solvent of ethanol or distilled water (Table 5). The diameter of the antimicrobial zone of the extract in three solvents of methanol, ethanol, or distilled water was 14.7 ± 0.91 mm; 12.7 ± 1.92 mm; 12.1 ± 1.59 mm, respectively, which was statistically higher than amoxicillin antibiotics (9.2 ± 5.3 mm) (Table 5). This demonstrated that the antimicrobial ability of *W. chinensis* extracts could be replaced by antibiotics in the prevention and treatment of diseases caused by *V. parahaemolyticus* in white-leg shrimp.

Based on the microbial resistance assessment table (Table 1), it is found that *W. chinensis* extracts in all three solvents (methanol, ethanol, or distilled water) had moderate antimicrobial effects toward *V. parahaemolyticus* (the diameter of the antimicrobial zone was 10–14 mm). Methanol is a highly polar solvent with a high capability of dissolving natural compounds in herbs, therefore, the extract of herbs in methanol showed the highest antimicrobial effect. The crude extract from the stem, and leaves of *Polygonum chinense* L. at doses of 10 $\mu\text{g/mL}$ showed the diameter of antibacterial zone against *V. parahaemolyticus* causing AHPND in brackish water shrimp reached to 20.6 mm [1]. Meanwhile, the extract of seed of *Rhodomyrtus tomentosa* at a concentration of 30 $\mu\text{g}/\mu\text{L}$ had the diameter of antibacterial zone toward *V. parahaemolyticus* KC13.14.2; *V. parahaemolyticus* KC12.02.0 and *Vibrio* sp. KC13.17.5 were 17.67, 18.00, and 19.33 mm, respectively. In addition, among seven types of herbal extracts, the extract of *Ricinus communis* L showed the highest efficacy antimicrobial on 2 strains of *V. harveyi* which caused luminous disease, and *V. parahaemolyticus* causing AHPND in shrimp, and its diameter of antimicrobial zone were 18.0 ± 1.4 mm and 17.5 ± 0.7 mm, respectively [4], [5]. Thus, the antagonistic of *W. chinensis* extract in methanol solvent obtained in this study to *V. parahaemolyticus* has moderate and strong antimicrobial capacity [5] reaching 14.7 ± 0.91 mm, however, this was smaller than crude extracts of herbs that used in the study of Hong Mong

Huyen et al (2020), but crude extracts are often more difficult to preserve than high extracts [4].

Results of the determination of the minimum inhibitory concentration (MIC) and the minimum bactericidal concentration (MBC) of *W. chinensis* extract for *V. parahaemolyticus* was presented in Table 6 and 7. In the extraction of *W. chinensis* in distilled water, bacteria still grow at the concentration of 1000 mg of extract/ L (OD = 0.2 after 24 hours of incubated bacteria with herb extract at 28-30 °C). The *W. chinensis* extract in methanol, or ethanol solvents inhibit the growth of *V. parahaemolyticus* at very low concentrations (15.5 mg/L).

However, bacteria re-grow on TSA plates after 24 hours of culture at concentrations of 15.5 mg/L for extract in methanol solvent and 31.25 mg/L for extract in ethanol solvent. From the results of screening and determination of minimum inhibitory concentration (MIC) on *V. parahaemolyticus* (Table 6), the minimum bactericidal concentration (MBC) of the extract on *V. parahaemolyticus* was presented in Table 7.

The MBC of *W. chinensis* extract on *V. parahaemolyticus* in distilled water or ethanol or methanol was 5000 mg/L, 62.5 mg/L, and 31.25 mg/L, respectively. According to Cadillac and Mourey (2001), if the MBC/MIC ratio is ≤ 4 , the herbal extract has been considered as a bacterial effect. If this ratio is greater than 4, they only inhibit the growth of bacteria [17]. From the results of the study, it is shown that the extract of *W. chinensis* in methanol, or ethanol solvents has a bacterial effect on *V. parahaemolyticus* causing acute hepatopancreatic necrosis AHPND on white leg shrimp with an MBC/MIC ratio was 2 and 4, respectively.

3.4. Results of the toxicity test

Based on the above experiment, the highest value of MBC of *W. chinensis* extract toward *V. parahaemolyticus* was used in this experiment for evaluating the toxicity of this extract in white-leg shrimp.

The result of the toxicity test showed the mortality occurred on the first day and finished on day 6 after feeding with feed mix with *W. chinensis* extract in treatments 2 and 3. After 7 days of feeding with *W. chinensis* extract at a concentration of 20 x MBC (625 mg/L) (treatment 2), the cumulative mortality was 16.7% and was lower than the mortality of shrimp in the treatment 3 (23.3%) which shrimp was fed with *W. chinensis* extract at a concentration of 40x MBC (1250 mg/L).

The highest cumulative mortality was observed in treatment 3 compared to other treatments. By contrast, no mortality was recorded during the experiment in treatment 4 (control) where shrimp fed with commercial feed without *W. chinensis* extract was added, and in treatment 1 which the shrimp fed with *W. chinensis* extract was added at a concentration of 1x MBC (31.25 mg/L) (Figure 9).

The cumulative mortality in all treatments was less than 50%, therefore the LD₅₀ has not been determined. No bacteria or none *V. parahaemolyticus* was recovered from the hepatopancreas of dead or moribund shrimp. However, dead and moribund shrimp showed pale liver (Figure 10). The *W. chinensis* contains biochemical compounds that have a high antibacterial and antifungal effect such as phenolic, saponin, alkaloid, and flavonoid, however, if the concentration of *W. chinensis* in feed was too high, this extract can hurt the liver of animal [7]. In this study, the dose of 625 mg/L or 1250 mg/L seemed too high for shrimp used, and the dose of 31.25 mg/L of *W. chinensis* was safe for shrimp.

4. Conclusions

- The extract of *W. chinensis* contains biochemical compounds such as fat, oil, alkaloid, carotenoid, flavonoid, organic acid, tannin, reduce agents
- The highest amount of polyphenol and flavonoid (74.33 mg GAE/ g extract and 24.59 mg RE/g extract, respectively), was identified in the *W. chinensis* extract in the methanol solvent followed by the 3.65 mg GAE/g extract (polyphenol) and 20.63 mg RE/g extract (flavonoid) were determined in the *W. chinensis* extract in the solvent of ethanol. The lowest total polyphenol and flavonoid (high 53.07 mg GAE/g and 3.20 mg RE/g extract, respectively) was observed in the *W. chinensis* extract in the distilled water solvent.
- The efficiency extraction of *W. chinensis* was 18.2% in a distilled water solvent, 11.3% in ethanol solvent, and 11% in methanol solvent.
- The extract of *W. chinensis* in all three solvents (distilled water, 99.8% methanol, and 96% ethanol) showed good antimicrobial effects toward *V. parahaemolyticus*, in which tract of *W. chinensis* in methanol solvent gave the best results with a diameter of antibacterial zone was 14.7 ± 0.91 mm.

Acknowledgments

The paper was funded by the Domestic Master/PhD scholarship programme of Vingroup Innovation Foundation (VINIF), Vingroup

Institute of big data (VINBIGDATA) [VINIF. 2022.TS.082]. This study was also partly supported by the Hue University Strong Research Group (NCM.DHH.2022.05) and by the Vietnam Ministry of Science and Technology (project code: ĐTĐLCN.56/22).

REFERENCES

1. Truong Thi My Hanh, Pham Thi Yen, Pham Thi Huyen, Huynh Thi My Le, Pham Thi Hong Minh, Do Tien Lam, Tran Thi Hoai Van, Phan Thi Van (2017), Antibacterialeffect of *Polygonum chinense* L. extract on pathogen bacteria of acute hepatopancreatic necrosis disease in brackish shrimps, *Vietnam Science and Technology Journal*, 17(6). (in Vietnamese)
https://b.vjst.vn/index.php/ban_b/article/download/486/477.
2. Trieu Thi Thanh Hang, Nguyen Cong Trang, Cao Tuan Duc and Le Thi Thuy Vy (2018). Study on the anti-bacterial activities of extracts from sakae naa (*Combretum quadrangulare*) on diseased aquatic animals-bacteria under *in vitro* conditions, *Can Tho University Scientific Journal*, 151-157. (in Vietnamese) doi: 10.22144/ctu.jsi.2018.048.5
3. Tran Hung, Nguyen Viet Kinh (2015), Research Methods for Medicinal Plants, University of Medicine and Pharmacy, Ho Chi Minh City, 126p. (in Vietnamese)
4. Hong Mong Huyen, Vo Tan Huy & Tran Thi Tuyet Hoa (2018), Antimicrobial activity of herbal extracts against shrimp pathogenic bacteria, *Can Tho University Science Journal*, 54(2), 143–150. (in Vietnamese)
<https://ctujsvn.ctu.edu.vn/index.php/ctujsvn/article/download/3178/641>.
5. Hong Mong Huyen, Le Quoc Viet, Tran Ngoc Hai, Tran Thi Tuyet Hoa (2020), Effects of herbal extracts on the growth, non-specific immune responses and disease resistance of white leg shrimp (*Penaeus vannamei*) against *Vibrio parahaemolyticus*, *Can Tho University Scientific Journal*, 56(5B), 150–159. (in Vietnamese)
<https://ctujsvn.ctu.edu.vn/index.php/ctujsvn/article/download/3605/214>.
6. Nguyen Quang Linh (2010), Application of Bokashi of betel leaf (*Piper betel*) for safe shrimp farming and the role of the community in Tam Giang - Cau Hai lagoon, Thua Thien Hue province. *Publish House of Hue University*; 2010.119 pp. (in Vietnamese)

7. Do Tat Loi (2000), *Vietnamese medicinal plants and herbs*, Publish House of Medicine, Ho Chi Minh city, Vietnam, 86pp. (in Vietnamese)
8. Tran Thanh Men, Nguyen Quoc Cuong, Nguyen Thi Anh Thu, Pham Lam Thao Quyen, Phan Cuc Phuong, Chiem Thi Ngoc Le, Nguyen Dinh Hai Yen and Do Tan Khang, (2019). Allelopathic effects of extracts from *Wedelia trilobata* (L.) Hitchc on seed germination. *Can Tho University Scientific Journal*, 55, 85 - 90. (in Vietnamese) doi:10.22144/ctu.jsi.2019.011
9. Nguyen Ngoc Phuoc, Nguyen Thi Xuan Hong, and Nguyen Cong Chung (2020), Study on the Virulence Factors of *Vibrio parahaemolyticus* Caused Acute Hepatopancreatic Necrosis Disease (AHPND) in Farmed White Leg Shrimp (*Litopenaeus vannamei*) in Thua Thien Hue Province, *Vietnam Journal of Agricultural Sciences*, 18(3), 202–211. (in Vietnamese) <http://tapchi.vnua.edu.vn/wp-content/uploads/2020/06/tap-chi-so-3.1.5-1.pdf>.
10. Nguyen Ngoc Phuoc., Nguyen Anh Tuan., Nguyen Quang Linh., Ngo Thi Huong Giang and Nguyen Nam Quang. (2007). Using herbs and herbal extracts in the treatment of bacterial diseases for aquatic animals. Proceedings of Science and Technology. *Vietnam Union of Science and Technology*. (in Vietnamese)
11. Bui Quang Te., & Le Xuan Thanh. (2006). Results of using (VTS1-C), and (VTS1-T), extracted from herbs for prevention and treatment of bacterial disease in black tiger shrimp and catfish. (in Vietnamese)
12. Trinh Thu Trang, Nguyen Thanh Hai (2016), In vitro bacteriostatic effect of dried betel nut extract (*Piper betel*) against *Aeromonas* spp. and *Streptococcus agalactiae* causing hemorrhagic disease in tilapia, *Vietnam Journal of Agricultural Sciences*, 14(6), 869–8761. (in Vietnamese)
13. Pham Thi Hai Yen, Nguyen Duy Quynh Tram, Hoang Ngoc Han, Tran Vinh Phuong (2019), Antibacterial activity of *Phyllanthus amarus* and effective microorganism (EM5) toward acute hepatopancreatic necrosis disease caused by *Vibrio parahaemolyticus* in white leg shrimps (*Litopenaeus vannamei*), *Hue University scientific journal: natural science*, 107–114. (in Vietnamese) <http://doi:10.26459/hueuni-jns.v128i1E.5440>
14. Ahn, K., (2017). The worldwide trend of using botanical drugs and strategies for developing global drugs. *BMB Reports*, 50 (3): 111–116. <https://doi.org/10.5483%2FBMBRep.2017.50.3.221>
15. Akiyama, H., Fujii, K., Yamasaki, O., Oono, T., and Iwatsuki, K. (2001). Antibacterial action of several tannins against *Staphylococcus aureus*. *Journal of antimicrobial chemotherapy*, 48(4): 487-491. <https://doi.org/10.1093/jac/48.4.487>
16. Blainski, A., Lopes, G.C., & de Mello, J. C. P. (2013), “Application and analysis of the folin-ciocalteu method for the determination of the total phenolic content from *limonium brasiliense* L.”, *molecules*, 18(6), pp. 6852-6865. <https://doi.org/10.3390/molecules18066852>
17. Cadillac, N., & Mourey, A. (2001), Antibacterial activity of the essential oil of *Picea excels* on *Listeria*, *Staphylococcus aureus* and coliform bacteria, *Food Microbiology*, 18(3), 261–268. <https://doi.org/10.1006/fmic.2000.0397>
18. Chakraborty, S.B., and Hancz, C. (2011). Application of phytochemicals as an immunostimulant, antipathogenic and antistress agent in finfish culture. *Reviews in Aquaculture*, 3(3): 103-119. <https://doi.org/10.1111/j.1753-5131.2011.01048>.
19. Citarasu, T., 2010. Herbal biomedicines: a new opportunity for the aquaculture industry. *Aquaculture International*, 18(3): 403-414 [https://doi.org/10.1016/0959-8030\(92\)90067-8](https://doi.org/10.1016/0959-8030(92)90067-8)
20. CLSI (1998), *Methods for Determining Bactericidal Activity of Antimicrobial Agents, Approved Guideline*, CLSI document M26-A, Clinical and Laboratory Standards Institute, 950 West Valley Roadn Suite 2500, Wayne, Pennsylvania 19087, USA. https://clsi.org/media/1462/m26a_sample.pdf.
21. CLSI (2012), *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically*, Approved Standard, 9th ed., CLSI document M07-A9, Clinical and Laboratory Standards Institute, 950 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087, USA. https://clsi.org/media/1632/m07a10_sample.pdf.
22. Cos P., Vlietinck A.J., Berghe D.V. & Maes L. (2006). The anti-infective potential of natural products: How to develop a stronger in vitro ‘proof of concept’. <https://doi.org/10.1016/j.jep.2006.04.003>
23. Courts F. L, and Williamsom G, 2015. The Occurrence, Fate and Biological of C-glycosyl Flavonoids in the Human Diet. *Critical*

- Reviews in Food Science and Nutrition, 55(10):1352-1367.
<https://doi.org/10.1080/10408398.2012.694497>
24. Doss, A., Mubarack, H.M., and Dhanabalan, R. (2009). Antibacterial activity of tannins from the leaves of *Solanum trilobatum* Linn. *International Journal of Systematic and Evolutionary Microbiology*, 2(2): 41-43. <http://www.indjst.org/>
25. Faikoh, E.N., Hong, Y.H., Hu, & S.Y. (2014), Liposome-encapsulated cinnamaldehyde enhances zebrafish (*Danio rerio*) immunity and survival when challenged with *Vibrio vulnificus* and *Streptococcus agalactiae*, *Fish & Shellfish Immunology*, 38, 15–24. <http://doi.org/10.1016/j.fsi.2014.02.024>.
26. Govindappa, M., Naga Sravya S., Poojashri M.N., et al., 2011. Antimicrobial, antioxidant, and in vitro anti-inflammatory activity and phytochemical screening of water extract of *Wedelia trilobata* (L.) Hitchc. *Journal of Medicinal Plants Research*. 5(24): 5718-5729 <https://doi.org/10.1006/fmic.2000.0397>.
27. Kumar, S., Narwal, S., Kumar, S., and Prakash, O., 2011. α -glucosidase inhibitors from plants: A natural approach to treat diabetes. *Pharmacognosy Review*, 5(9): 19-29. <http://doi.org/10.4103/0973-7847.79096>
28. Makuasa D. A., Ningsih P. (2020), "The Analysis of Total Flavonoid Levels in Young Leaves and Old Soursop Leaves (*Annona muricata* L.) Using UV-Vis Spectrophotometry Methods", *Journal of Applied Science, Engineering, Technology, and Education*, 2(1), pp. 11-17. <https://doi.org/10.35877/454RI.asci2133>
29. Milan V., Bharat M. (2019), "Qualitative analysis, total phenol content (TPC) and total tannin content (TTC) by using different solvent for flower of *Butea monosperma* (Lam.) Taub. collected from Saurashtra region", *Journal of Pharmacognosy and Phytochemistry*, 8(3), pp. 2902-2906. <https://www.phytojournal.com/archives/2019/vol8issue3/PartAO/8-2-615-581.pdf>
30. Muhammad, A., and Mudi, Y. (2011). Phytochemical screening and antimicrobial activities of *Terminalia catappa*, leaf extracts. *Biokemistri*, 23(1): 35-39. <https://www.ajol.info/index.php/biokem/article/view/77668>
31. Munaeni, W., Yuhana, M., Setiawati, M., and Wahyudi, A.T. (2019). Phytochemical analysis and antibacterial activities of *Eleutherine bulbosa* (Mill.) Urb. extract against *Vibrio parahaemolyticus*. *Asian Pacific Journal of Tropical Biomedicine*, 9(9): 397. <http://doi.org/10.4103/2221-1691.267669>
32. Neelam B., Nadpi G.K, Titpawan N., Kholeeyoh J., and Teerapol S., 2012. Evaluation of the wound healing potential of *Wedelia trilobata* (L.) leaves. *Journal of Ethnopharmacology*. 141: 817–824. <https://doi.org/10.1016/j.jep.2012.03.019>
33. Phạm Thi Thu Ha, Nguyen Thi Bao Tran, Nguyen Thi Ngọc Tram, Vo Hoang Kha (2020), "Total phenolic, total flavonoid contents and antioxidant potential of Common Bean (*Phaseolus vulgaris* L.) in Vietnam", *AIMS Agriculture and Food*, 5(4), pp. 635-648. DOI: 10.3934/agrfood.2020.4.635
34. Reed, L. J., & Muench, H. (1938). A simple method of estimating fifty percent endpoints. *American Journal of Hygiene* 27(3), 493-497.) <https://doi.org/10.1093/oxfordjournals.aje.a118408>
35. Turker, H., Yıldırım, A.B. and Karakaş, F.P. (2009), Sensitivity of Bacteria Isolated from Fish to Some Medicinal Plants, *Turkish Journal of Fisheries and Aquatic Sciences*, 9, 181–186. <https://dergipark.org.tr/en/pub/trjfas-ayrildi/issue/13279/1604>

